



Order: (888)-282-5810 (Phone)
(818)-707-0392 (Fax)
order@abiocode.com
Web: www.Abiocode.com

CXCR4 (C) Antibody, Rabbit Polyclonal

Cat#: R0067-2

Quantity: 100 ul

Predicted M.W.: 40 kDa

Lot#: Refer to vial

Application: WB

Uniprot ID: P61073

Background:

Chemokine (C-X-C motif) receptor 4 (CXCR4) is the receptor for cell-derived factor-1. It transduces signals that increase intracellular calcium ion levels. It also enhances MAPK/MAPK3 activation and reduces cellular cAMP levels. It is involved in hematopoiesis and in cardiac ventricular septum formation. CXCR4 also plays a role in vascularization of the gastrointestinal tract. Mutations in the gene that codes for CXCR4 have been associated with WHIM (warts, hypogammaglobulinemia, infections and myelokathexis) syndrome. CXCR4 acts as a co-receptor, along with CD4 protein to support HIV entry into cells. CXCR4 is also highly expressed in breast cancer cells.

Other Names:

CD184, D2S201E, FB22, HM89, HSY3RR, LAP3, LCR1, LESTR, NPY3R, NPYR, NPYRL, NP

Source and Purity:

Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing the C-terminal region of human CXCR4. Antibodies were purified by affinity purification using immunogen.

Storage Buffer and Condition:

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN₃. Store at -20 °C. Stable for 6 months from date of receipt.

Species Specificity:

Human

Tested Applications:

WB: 1:1,000-1:5,000 (detect endogenous protein*)

*: The apparent protein size on WB may be different from the calculated M.W. due to modifications.

Product Data:




Fig 1. Western blot of total cell extracts from human Jurkat, using Ab (R0067-2) at RT for 2 h.