



Order: (888)-282-5810 (Phone)
(818)-707-0392 (Fax)
order@abiocode.com
Web: www.Abiocode.com

CCL17 (N) Antibody, Rabbit Polyclonal

Cat#: R0132-1

Quantity: 100 ug

Predicted M.W.: 11 kDa

Lot#: Refer to vial

Application: ELISA

Uniprot ID: Q92583

Background:

Chemokine (C-C motif) ligand 17 (CCL17) is a member of the subfamily of small cytokine CC genes characterized by two adjacent cysteine residues. CCL17 is clustered on the q arm of chromosome 16. It has been shown to display chemotactic activity for T lymphocytes but not monocytes or granulocytes. CCL17 has the ability to bind to CCR4 and CCR8. CCL17 plays a role in T-cell development in thymus along with trafficking and activation of mature T cells.

Other Names:

CC chemokine TARC, Small-inducible cytokine A17, Thymus and activation-regulated chemokine, Scya17, Scya17l, Abcd-2, Tarc

Source and Purity:

Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing the N-terminal region of mouse CCL17. Antibodies were purified by affinity purification using immunogen.

Storage Buffer and Condition:

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN₃. Store at -20 °C. Stable for 6 months from date of receipt.

Species Specificity:

Mouse

Tested Applications:

ELISA: 1:1,000- 1:10,000 (detect native recombinant protein in a direct ELISA assay).
Suitable for using as ELISA capture or detection antibodies.

Last Update: 03/2011

**For research use only. Not for therapeutic or diagnostic purposes.
Abiocode, Inc., 29397 Agoura Rd., Ste 106, Agoura Hills, CA 91301**