

NLK (C) Antibody, Rabbit Polyclonal

Cat#: R0838-2 Quantity: 100 ul Predicted I Observed M.W.: 58 I 65 kDa Lot#: Refer to vial Application: WB Uniprot ID: Q9UBE8

Background:

NLK regulates a number of transcription factors with key roles in cell fate determination. It is a positive effector of the non-canonical Wnt signaling pathway, acting downstream of WNT5A, MAP3K7/TAK1 and HIPK2.

Other Names:

LAK1

Source and Purity:

Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing C-terminal region of human NLK. Antibodies were purified by affinity purification using immunogen.

Storage Buffer and Condition:

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN₃. Store at -20 °C. Stable for 6 months from date of receipt.

Species Specificity:

Human, Mouse

Tested Applications:

WB: 1:1,000-1:3,000 (detect endogenous protein*)

*: The apparent protein size on WB may be different from the calculated M.W. due to modifications.



Product Data:

.




Fig 1. Western blot of total cell extracts from mouse brain; using anti-NLK (C) (R0838-2) at RT for 2 h.

Last Update: 11/2012