



Order: (888)-282-5810 (Phone)
(818)-707-0392 (Fax)
order@abiocode.com
Web: www.Abiocode.com

RLIM (N) Antibody, Rabbit Polyclonal

Cat#: R2807-1

Quantity: 100 ul

Predicted | Observed MW: 69 kDa

Lot#: Refer to vial

Application: WB

Uniprot ID: Q9NVW2

Background:

RING finger LIM domain-binding protein (RLIM) is a RING-H2 zinc finger protein. It has been shown to be an E3 ubiquitin protein ligase that targets LIM domain binding 1 (LDB1/CLIM), and causes proteasome-dependent degradation of LDB1. RLIM and LDB1 are co-repressors of LHX1/LIM-1, a homeodomain transcription factor [provided by RefSeq].

Other Names:

E3 ubiquitin-protein ligase RLIM, LIM domain-interacting RING finger protein, RING finger LIM domain-binding protein, R-LIM, RING finger protein 12, Renal carcinoma antigen NY-REN-43, RNF12, MGC15161

Source and Purity:

Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing the N-terminal region of human RLIM. Antibodies were purified by affinity purification using immunogen.

Storage Buffer and Condition:

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN₃. Store at -20 °C. Stable for 6 months from date of receipt.

Species Specificity:

Human

Tested Applications:

WB: 1:1,000-1:3,000 (detect endogenous protein*)

*: The apparent protein size on WB may be different from the calculated M.W. due to modifications.

Product Data:




Fig 1. Western blot of total cell extracts from human HeLa, using anti- RLIM (N) (R2807-1) at RT for 2 h.