

Order: (888)-282-5810 (Phone) (818)-707-0392 (Fax) order@abiocode.com Web: www.Abiocode.com

TET1 (N2) Antibody, Rabbit Polyclonal

Cat#: R1084-1 Quantity: 100 ul Predicted M.W.: 235 kDa Lot#: Refer to vial Application: WB Uniprot ID: Q8NFU7

Background:

TET1 is a methylcytosine dioxygenase that catalyzes the conversion of the modified genomic base 5methylcytosine (5mC) into 5-hydroxymethylcytosine (5hmC). Methylation at the C5 position of cytosine bases is an epigenetic modification of the mammalian genome which plays an important role in transcriptional regulation. TET1 might initiate a process leading to cytosine demethylation through deamination into 5-hydroxymethyluracil (5hmU) and subsequent replacement by unmethylated cytosine by the base excision repair system. TET1 preferentially binds to CpG-rich sequences at promoters of both transcriptionally active and polycomb-repressed genes. By controlling the levels of 5mC and 5hmC at gene promoters, it may regulate the gene silencing induced by cytosine methylation. It has been shown that TET1 plays an important role in embryonic stem cells maintenance and regulating the balance between pluripotency and lineage commitment.

Other Names:

Methylcytosine dioxygenase TET1, CXXC-type zinc finger protein 6, Leukemia-associated protein with a CXXC domain, Ten-eleven translocation 1 gene protein, CXXC6, KIAA1676, LCX

Source and Purity:

Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing the N-terminal region of human TET1. Antibodies were purified by affinity purification using immunogen.

Storage Buffer and Condition:

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN₃. Store at -20 °C. Stable for 6 months from date of receipt.

Species Specificity:

Human, Mouse

Tested Applications:

WB: 1:500-1:2,000

*: The apparent protein size on WB may be different from the calculated M.W. due to modifications.



Product Data:




Fig 1. Western blot of total cell extracts from human 293 T cells transfected with the empty vector, or Flag-tagged murine TET1, TET2 or TET3; using indicated Abs at RT for 2 h.