



Order: (888)-282-5810 (Phone)  
(818)-707-0392 (Fax)  
[order@abiocode.com](mailto:order@abiocode.com)  
Web: [www.Abiocode.com](http://www.Abiocode.com)

---

## **RUNX1T1 (C) Antibody, Rabbit Polyclonal**

**Cat#: R2572-2**

**Quantity: 100 ul**

**Predicted | Observed M.W.: 68 | 75 kDa**

**Lot#: Refer to vial**

**Application: WB**

**Uniprot ID: Q06455**

### **Background:**

Runt-related transcription factor 1 (RUNX1T1) is a member of the myeloid translocation gene family, which interact with DNA-bound transcription factors and recruit a range of co-repressors to facilitate transcriptional repression. The t(8;21)(q22;q22) translocation is one of the most frequent karyotypic abnormalities in acute myeloid leukemia. The translocation produces a chimeric gene made up of the 5'-region of the RUNX1T1 gene fused to the 3'-region of this gene. The chimeric protein is thought to associate with the nuclear corepressor/histone deacetylase complex to block hematopoietic differentiation. Alternative splicing of the RUNX1T1 gene results in multiple transcript variants [provided by RefSeq].

### **Other Names:**

Runt-related transcription factor 1, Protein CBFA2T1, Cyclin-D-related protein, Eighty two one protein, Protein ETO, Protein MTG8, Zinc finger MYND domain-containing protein 2, AML1T1, CBFA2T1, CDR, ETO, MTG8, ZMYND2

### **Source and Purity:**

Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing the C-terminal region of human RUNX1T1. Antibodies were purified by affinity purification using immunogen.

### **Storage Buffer and Condition:**

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN<sub>3</sub>. Store at -20 °C. Stable for 6 months from date of receipt.

### **Species Specificity:**


Human, Mouse

### **Tested Applications:**

WB: 1:1,000-1:3,000 (detect endogenous protein\*)

\*: The apparent protein size on WB may be different from the calculated M.W. due to modifications.

**Product Data:**



**Fig 1.** Western blot of total cell extracts from A, C) mouse brain, B, D) mouse thymus; using 2 independent Abs against 2 distinct regions of human RUNX1T1 at RT for 2 h.