



Order: (888)-282-5810 (Phone)
(818)-707-0392 (Fax)
order@abiocode.com
Web: www.Abiocode.com

NCAM1 (vPair™) Antibodies

Cat#: R2444-vp

Lot#: Refer to vial

Predicted | Observed M.W.: 95 | 140 kDa

Uniprot ID: P13591

Application: WB

Quantity: 50 ul NCAM1 (N) (R2444-1) Rabbit Polyclonal Antibody &
50 ul NCAM1 (C) (R2444-2) Rabbit Polyclonal Antibody

Product Introduction:

vPair™ antibodies represent a pair of fully characterized antibodies that recognize two different regions of a target protein. The product is developed by Abiocode to address whether the signal observed truly represents the protein of interest, an often encountered issue in antibody-based assays. The use of a pair of fully characterized vPair™ antibodies in the same assay can validate signal specificity since vPair™ antibodies recognize two independent epitopes of the same protein. Different sets of vPair™ antibodies are developed at Abiocode to work with specific applications, including antibody arrays, Western blot, IP-Western, CHIP, IHC, and FACS.

Background:

NCAM1 is involved in cell-to-cell interactions as well as cell-matrix interactions during development and differentiation. NCAM1 has been shown to be involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance.

Other Names:

Neural cell adhesion molecule 1, N-CAM-1, NCAM-1, CD_antigen=CD56, NCAM

Source and Purity:

Rabbit polyclonal antibodies were produced by immunizing animals with GST-fusion proteins containing either N-terminal [NCAM1 (N) (R2444-1)] or the C-terminal [NCAM1 (C) (R2444-2)] region of human NCAM1. Antibodies were purified by affinity purification using immunogen.

Storage Buffer and Condition:

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN₃. Store at -20 °C. Stable for 6 months from date of receipt.

Species Specificity:

Human, Mouse

Tested Applications:

WB: 1:1,000-1:3,000 (detect endogenous protein*)

*: The apparent protein size on WB may be different from the calculated M.W. due to modifications.

Product Data:




Fig 1. Western blot of total cell extracts from A, B) mouse brain, using 2 independent Abs against 2 distinct regions of human NCAM1 at RT for 2 h. These 2 Abs appear to recognize the 140 kD form of NCAM1.