

PIF3 (M) Antibody, Rabbit Polyclonal

Cat#: R3392-3 Quantity: 100 ul Predicted I Observed M.W.: 57 I 63kDa Lot#: Refer to vial Application: WB Uniprot ID: O80536

Background:

PIF3 is a transcription factor acting positively in the phytochrome signaling pathway. PIF3 activates transcription by binding to the G box (5'-CACGTG-3').

Other Names:

Transcription factor PIF3, Basic helix-loop-helix protein 8, AtbHLH8, bHLH 8, Phytochromeassociated protein 3, Phytochrome-interacting factor 3, Transcription factor EN 100, bHLH transcription factor bHLH008, BHLH8, EN100, PAP3, At1g09530, F14J9.19

Source and Purity:

Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing the middle region of *arabidopsis thaliana* PIF3 (AT1G09530). Antibodies were purified by affinity purification using immunogen.

Storage Buffer and Condition:

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN₃. Store at -20 °C. Stable for 6 months from date of receipt.

Species Specificity:

Arabidopsis thaliana

Tested Applications:

WB: 1:500-1:2,000 (detect endogenous protein*)

*: The apparent protein size on WB may be different from the calculated M.W. due to modifications.



Product Data:




Fig 1. A) Western blot of total protein extracts from human 293T cells transfected with Flag-tagged Arabidopsis PIF1, Myc-tagged Arabidopsis PIF3, PIF4 or PIF5, using a mixture of anti-Flag and anti-Myc Abs or anti-PIF3 (M) (R3392-3) Ab as indicated at RT for 2h. **B**) Same as **A** except that protein samples were prepared from wild type (WT) or pifq mutant (Mut, PIF1/3/4/5–deficient mutant) grown in dark for 4 days; using anti-PIF3 (M) (R3392-3) at RT for 2 h. All protein samples were prepared using 4xSDS sample buffer.