



Order: (888)-282-5810 (Phone)  
(818)-707-0392 (Fax)  
[order@abiocode.com](mailto:order@abiocode.com)  
Web: [www.Abiocode.com](http://www.Abiocode.com)

---

## **TET1 (C) Antibody, Rabbit Polyclonal**

**Cat#: R1084-5**

**Quantity: 100 ul**

**Predicted | Observed M.W.: 219 | 250 kDa**

**Lot#: Refer to vial**

**Application: WB**

**Uniprot ID: Q3URK3**

### **Background:**

TET1 is a methylcytosine dioxygenase that catalyzes the conversion of the modified genomic base 5-methylcytosine (5mC) into 5-hydroxymethylcytosine (5hmC). Methylation at the C5 position of cytosine bases is an epigenetic modification of the mammalian genome which plays an important role in transcriptional regulation. TET1 might initiate a process leading to cytosine demethylation through deamination into 5-hydroxymethyluracil (5hmU) and subsequent replacement by unmethylated cytosine by the base excision repair system. TET1 preferentially binds to CpG-rich sequences at promoters of both transcriptionally active and polycomb-repressed genes. By controlling the levels of 5mC and 5hmC at gene promoters, it may regulate the gene silencing induced by cytosine methylation. It has been shown that TET1 plays an important role in embryonic stem cells maintenance and regulating the balance between pluripotency and lineage commitment.

### **Other Names:**

Methylcytosine dioxygenase TET1, CXXC-type zinc finger protein 6, Leukemia-associated protein with a CXXC domain, Ten-eleven translocation 1 gene protein, CXXC6, KIAA1676, LCX

### **Source and Purity:**

Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing the C-terminal region of mouse TET1. Antibodies were affinity-purified using the immunogen.

### **Storage Buffer and Condition:**

Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN<sub>3</sub>. Store at -20 °C. Stable for 6 months from date of receipt.

### **Species Specificity:**

Mouse

### **Tested Applications:**


WB: 1:1,000-1:3,000

\*: The apparent protein size on WB may be different from the calculated M.W. due to modifications.

---

**For research use only. Not for therapeutic or diagnostic purposes.  
Abiocode, Inc., 29397 Agoura Rd., Ste 106, Agoura Hills, CA 91301**

**Product Data:**



**Fig 1. A)** Western blot of total cell extracts from human 293T cells transfected with a vector control or Flag-tagged mouse TET1 (Flag-TET1), using anti-Flag or anti-TET1 (C) (R1084-5) at RT for 2 h.